


Motivation


Given a bipartite graph, is it the canonical double cover (CDC) of some base graph?



• Are there non-isomorphic base graphs that share a given CDC?

Motivation


Given a bipartite graph, is it the canonical double cover (CDC) of some base graph?



• Are there non-isomorphic base graphs that share a given CDC?

Motivation


Given a bipartite graph, is it the canonical double cover (CDC) of some base graph?



• Are there non-isomorphic base graphs that share a given CDC?

Motivation

Given a bipartite graph, is it the canonical double cover (CDC) of some base graph?



• Are there non-isomorphic base graphs that share a given CDC?


Factorization under direct product

Factorization under direct product

W. Imrich and T. Pisanski (2008): The factorization of a non-bipartite connected graph under direct product is unique.

R. Hammack, W. Imrich and S. Klavzar (2011): For a bipartite connected graph, factorization may not be unique.

D. Witte Morris (2022): Bip Conn Gr has a bipartite factor which may NOT be unique.



The CDC is Bipartite

The canonical double cover (CDC) of a base graph G

is its direct product $G \times K_2$ with $K_2 = a \longleftrightarrow b$.

Edge $(u, a), (w, b)$ is an edge of CDC iff
 (u, w) is an edge of G and (a, b) is the edge of K_2 .

Also if a graph is a CDC, then its bipartite factor under direct product is NOT necessarily K_2 .

The CDC is Bipartite


The canonical double cover (CDC) of a base graph G

is its direct product $G \times K_2$ with $K_2 = a \longleftrightarrow b$.


Edge $(u, a), (w, b)$ is an edge of CDC iff
 (u, w) is an edge of G and (a, b) is the edge of K_2 .

Also if a graph is a CDC, then its bipartite factor under direct product is NOT necessarily K_2 .


Construction of a CDC



G




H




CDC

No labelling of H gives the same labelling of its CDC as that of G.


Construction of a CDC



G



H



CDC

No labelling of H gives the same labelling of its CDC as that of G.

Inverse Problem

Given a bipartite connected graph B

1. Is it a CDC?
2. Determine base graphs G such that $B = CDC(G)$.

The problem is intractable.

1. The **complexity** of algorithms known to date is factorial, $O(\frac{n}{2})!$
2. **Open Problem:** Is it fixed parameter tractable?

Inverse Problem

Given a bipartite connected graph B

1. Is it a CDC?
2. Determine base graphs G such that $B = CDC(G)$.

The problem is intractable.

1. The **complexity** of algorithms known to date is factorial, $O(\frac{n}{2})!$
2. **Open Problem:** Is it fixed parameter tractable?

Existence of a base graph

Starting from the adjacency matrix

$$\mathbf{A}(B) = \left(\begin{array}{c|c} \mathbf{O} & \mathbf{K} \\ \hline \mathbf{K}^\top & \mathbf{O} \end{array} \right), \text{ of a bipartite graph } B,$$

if there exists a graph G with adjacency matrix permutationally equivalent to \mathbf{K} and \mathbf{K}^\top , then $B = \text{CDC}(G)$.

Sufficient condition

If there exist permutation matrices P and Q such that $P^\top \mathbf{K} Q$ is a real 0–1–symmetric matrix $\mathbf{A}(G)$, then B is the CDC of G in Γ_0 .

$$\mathbf{A}(\text{CDC}) = \left(\begin{array}{c|c} \mathbf{O} & \mathbf{A}(G) \\ \hline \mathbf{A}(G) & \mathbf{O} \end{array} \right),$$

by permuting the rows and columns of \mathbf{K} in different ways.

Existence of a base graph

Starting from the adjacency matrix

$$\mathbf{A}(B) = \left(\begin{array}{c|c} \mathbf{O} & \mathbf{K} \\ \hline \mathbf{K}^\top & \mathbf{O} \end{array} \right), \text{ of a bipartite graph } B,$$

if there exists a graph G with adjacency matrix permutationally equivalent to \mathbf{K} and \mathbf{K}^\top , then $B = \text{CDC}(G)$.

Sufficient condition

If there exist permutation matrices P and Q such that $P^\top \mathbf{K} Q$ is a real 0–1–symmetric matrix $\mathbf{A}(G)$, then B is the CDC of G in Γ_0 .



$$\mathbf{A}(\text{CDC}) = \left(\begin{array}{c|c} \mathbf{O} & \mathbf{A}(G) \\ \hline \mathbf{A}(G) & \mathbf{O} \end{array} \right),$$

by permuting the rows and columns of \mathbf{K} in different ways.

L.Porcu(1976), T.Pisanski, D.Marûsić, D.Surowski, W.Imrich,
R. Hammack, S. Klavzar, S. Wilson, R. Scapellato, N.
ZagagliaS, W. Pacco, J. Lauri, R. Mizzi, L. Collins, I.Sciriha

Unstable base graphs arise from Unexpected Symmetries



in the automorphism group of a CDC. $\text{Aut}(\text{CDC})$ contains group $\text{Aut}(G) \times \text{Aut}(K_2)$. If inclusion is strict, there may be symmetries in $\text{Aut}(\text{CDC})$ NOT lifted from $\text{AUT}(G)$.



L.Porcu(1976), T.Pisanski, D.Marûsić, D.Surowski, W.Imrich,
R. Hammack, S. Klavzar, S. Wilson, R. Scapellato, N.
ZagagliaS, W. Pacco, J. Lauri, R. Mizzi, L. Collins, I.Sciriha

Unstable base graphs arise from Unexpected Symmetries

in the automorphism group of a CDC. $\text{Aut}(\text{CDC})$ contains group $\text{Aut}(G) \times \text{Aut}(K_2)$. If inclusion is strict, there may be symmetries in $\text{Aut}(\text{CDC})$ NOT lifted from $\text{AUT}(G)$.



Unstable base graphs

Given a bipartite connected graph with no duplicate vertices isomorphic to CDC $G \times K_2$,
are there other base graphs H permutationally equivalent to G ?

Reduction of the Inverse Problem to

Determining all 2–1 surjective projection mappings from the vertices of the given CDC to the vertices of a base graph, respecting adjacencies.

As 2 vertices of the CDC are identified to obtain G' ,
new automorphisms may appear that were not in $\text{aut}(\text{CDC})$.

Unstable base graphs





Given a bipartite connected graph with no duplicate vertices isomorphic to CDC $G \times K_2$, are there other base graphs H permutationally equivalent to G ?

Reduction of the Inverse Problem to

Determining all 2–1 surjective projection mappings from the vertices of the given CDC to the vertices of a base graph, respecting adjacencies.






As 2 vertices of the CDC are identified to obtain G' , new automorphisms may appear that were not in $\text{aut}(\text{CDC})$.

Existence of Unstable Base Graphs



$$G_1 \cdot P_{12} \cdot P_{56} \cdot P_{34} = H$$

Existence of Unstable Base Graphs



$$G_1 \cdot P_{12} \cdot P_{56} \cdot P_{34} = H$$

Permutationally Equivalent Base Graphs

Collins and Sciriha (2020)

Base graphs with isomorphic CDCs have the same k -walk matrix $\mathbf{W}_G(k) = \left(\mathbf{j} \mid \mathbf{A}\mathbf{j} \mid \mathbf{A}^2\mathbf{j} \mid \dots \mid \mathbf{A}^{k-1}\mathbf{j} \right)$. Vertices corresponding to equal rows of $\mathbf{W}_G(k)$ lie in one part of the WALK vertex partition and are assigned one colour in the **WALK COLOURING**.

Permutationally Equivalent Base Graphs

Collins and Sciriha (2020)


Base graphs with isomorphic CDCs have the same k -walk matrix $\mathbf{W}_G(k) = \left(\mathbf{j} \mid \mathbf{A}\mathbf{j} \mid \mathbf{A}^2\mathbf{j} \mid \dots \mid \mathbf{A}^{k-1}\mathbf{j} \right)$. Vertices corresponding to equal rows of $\mathbf{W}_G(k)$ lie in one part of the WALK vertex partition and are assigned one colour in the **WALK COLOURING**.

Degree Sequence of Base Graphs

Given by the 2nd column of $\mathbf{W}_G(k)$

- Two graphs with isomorphic CDCs have the same degree sequence.
- Graphs have the same degree sequence iff they are Ryser-switch transformable.

A Ryser-Switch on two disjoint edges




Degree Sequence of Base Graphs

Given by the 2nd column of $\mathbf{W}_G(k)$

- Two graphs with isomorphic CDCs have the same degree sequence.
- Graphs have the same degree sequence iff they are Ryser-switch transformable.

A Ryser-Switch on two disjoint edges



Construction of an unstable base graphs from a stable one

Theorem

Let G and H be graphs having isomorphic CDCs. Then H is isomorphic to a graph obtained from G by RS of disjoint edges each with two common but distinct walk-coloured end-vertices.

Base graphs G and H have the same degree sequence.

So there exist a sequence of Ryser switches involving pairs of disjoint edges $\{u, v\}$ and $\{x, y\}$ in G with the same walk-colouring that are transformed to $\{u, x\}$ and $\{v, y\}$ in H , retaining the walk-colouring, leaving all other adjacencies intact.

Equivalent to permuting the rows and columns of G differently.

Construction of an unstable base graphs from a stable one

Theorem

Let G and H be graphs having isomorphic CDCs. Then H is isomorphic to a graph obtained from G by RS of disjoint edges each with two common but distinct walk-coloured end-vertices.


Base graphs G and H have the same degree sequence.

So there exist a sequence of Ryser switches involving pairs of disjoint edges $\{u, v\}$ and $\{x, y\}$ in G with the same walk-colouring that are transformed to $\{u, x\}$ and $\{v, y\}$ in H , retaining the walk-colouring, leaving all other adjacencies intact.

Equivalent to permuting the rows and columns of G differently.


The Converse is False

Not every RS on disjoint edges with the same walk-colouring gives non-isomorphic graphs with the same CDC.





The Converse is False


Not every RS on disjoint edges with the same walk-colouring gives non-isomorphic graphs with the same CDC.




Construction of infinite families of pairs of base graphs with the same CDC



Construction of infinite families of pairs of base graphs with the same CDC



Construction of infinite families of pairs of base graphs with the same CDC



Website

<https://maths.mt/walks/same-cdc/>